

## PLESIOGAMY, A TERM CONTRASTING WITH HERKOGAMY

GUY L. NESOM

2925 Hartwood Drive  
Fort Worth, Texas 76109  
guynesom@sbcglobal.net

### ABSTRACT

Plesiogamy is proposed for use as the term referring to a close positioning of stigma and anthers, which usually leads to autogamy. Herkogamy, the contrasting term, refers to a spatial separation of the stigma and anthers, thus increasing the likelihood of outcrossing.

**KEY WORDS:** herkogamy and allogamy, plesiogamy and autogamy, dichogamy, synchronogamy

**Herkogamy** (Greek, *herkos*, wall or fence, and *gamos*, marriage; Brown 1956) refers to a spatial separation of the stigma and anthers, thus increasing the likelihood of outcrossing and production of genetically variable offspring. In "approach herkogamy," the most common form of herkogamy, the stigma is above the level of the anthers; in "reverse herkogamy" the stigma is below the anthers.

No term appears to exist, however, in reference to the contrasting situation, where stamens and stigma of a single flower are at the same level, in essentially the same position, thus making it likely that self-pollination occurs. The term **plesiogamy** (Greek, *plesios*, near, alluding to the proximity of the stamens and stigma) is proposed here for this arrangement and is used in several studies dealing with the taxonomy of *Mimulus* sensu lato (e.g., Nesom 2012). Plesiomorphic refers to an unspecialized condition (in contrast to plesiogamy, which usually is interpreted as a specialization), but the word itself (Greek, *plesios*, near, *morphe*, form) intends to convey the sense of nearness to the original morphological form, i.e., primitive.

Perhaps "non-herkogamy" would serve just as well as a contrasting term to herkogamy, but a positive term (vs. one that simply negates the opposing condition) seems useful. We prefer electron to "non-proton," past to "non-future," yang to "non-yin," etc, and the contrasting autogamy/allogamy and chasmogamy/cleistogamy have long been in use.

**Autogamy** (self-pollination, from anthers to stigma of a single individual, commonly within the same flower) is the mode implied by plesiogamous arrangement of anthers and stigma. **Allogamy** (cross-pollination, pollination between flowers of different individuals; synonyms or near-synonyms exogamy, xenogamy) is the usual mode of pollination and fertilization implied by herkogamous arrangement of anthers and stigma.

**Dichogamy** (in plants) refers to a temporal separation of pollen shedding and stigma receptivity. Protandry and protogyny describe conditions when anthers or stigmas mature first. Apparently no contrasting term exists for dichogamy, and it's suggested that, if necessary or desirable, **synchronogamy** (Greek, *syn*, same or together, *chronos*, time) could be used to indicate that pollen shedding and stigma receptivity occur simultaneously.

To complete the brief survey of opposing "-gamy" terms, **chasmogamy** refers to the condition in which corollas are open at anthesis, exposing the anthers and stigma and usually leading



to outcrossing. **Cleistogamy** refers to corollas that are closed at anthesis, strongly correlated with self-pollination.

**Geitonogamy** (Greek, *geiton*, neighbor) is another related term. It refers to the pollination — by insects or by wind — of a flower with the pollen from another flower on the same plant. The result of geitonogamy is genetically similar to autogamy.

#### ACKNOWLEDGEMENTS

Many thanks to Beryl Simpson, Steve Weller, and Michelle Dudash for helpful comments.

#### LITERATURE CITED

- Brown, R.W. 1956. *Composition of Scientific Words* (rev. ed.). Smithsonian Institution Press, Washington, DC.
- Nesom, G.L. 2012. Taxonomy of *Erythranthe* sect. *Simiola* (Phrymaceae) in the USA and Mexico. *Phytoneuron* 2012-40: 1–123.